Supplementary Table 1. Preoperative and postoperative contraceptive preferences (McNemar analysis)					
Contraceptive methods	**Preoperative (n)**	**Postoperative (n)**	**McNemar (exact)**	**p**	**Bonferroni-corrected p-value**
None	66	49	p <0.001	**0.000104**	**0.000624**
BTL/BS	12	25	p <0.001	**0.000874**	**0.005244**
IUD	26	25	-	1.000000	1.000000
Withdrawal	9	6	-	0.248213	1.000000
Condom	3	4	-	1.000000	1.000000
Oral contraceptive pills	3	4	-	1.000000	1.000000
Comparison of contraceptive methods used in the preoperative and postoperative periods among 122 patients. McNemar's test was applied to assess statistical differences in the distribution of each method. Rare or clinically distinct methods were excluded from the analysis for clarity and consistency BTL: Bilateral tubal ligation, BS: Bilateral salpingectomy, IUD: Intrauterine device					

Supplementary Table 2. Summary of 25 patients undergoing bilateral tubal ligation or bilateral salpingectomy			
Variable	**Number of patients (n)**	**Percentage (%)**	**p**
Preoperative BTL/BS use	11	44.0	
Preoperative-other or no method	14	56.0	
Postoperative BTL/BS use	25	100.0	**0.031**
McNemar's exact test was applied to assess the within-subject difference BTL: Bilateral tubal ligation, BS: Bilateral salpingectomy			