



# Permanent Junctional Reciprocating Tachycardia-induced Dilated Cardiomyopathy: A Case Report

## *Permanent Resiprokan Kavşak Taşikardisine Bağlı Dilate Kardiyomiyopati: Bir Olgu Sunumu*

Fatih Battal, Şule Yıldırım, Hakan Aylanç, Fatih Köksal Binnetoğlu, Nazan Kaymaz, Celal Akdeniz\*

Çanakkale Onsekiz Mart University Faculty of Medicine, Department of Pediatrics, Çanakkale, Turkey

\*İstanbul Medipol University Faculty of Medicine, Department of Pediatrics, İstanbul, Turkey

### Abstract

We present a four-year-old girl who was admitted to our hospital with the complaints of dyspnea, tachypnea, cough, excess sweating and fatigue. Electrocardiogram (ECG) in the tachycardic girl showed inverted P waves in leads 2, 3 and aVF along with a P-R interval of 0.16 sec and an R-P interval of 0.28 sec. Transthoracic echocardiography revealed an enlarged and spherical left ventricle with diminished systolic functions. Holter ECG confirmed long R-P tachycardia with a rate of 140-160 beats/minute. She was diagnosed as having permanent junctional reciprocating tachycardia-induced dilated cardiomyopathy and successfully treated with catheter ablation and flecainide.

**Keywords:** Electrocardiography, child, catheter ablation

### Öz

Bu olgu sunumunda hastanemize dispne, takipne, öksürük, aşırı terleme ve yorgunluk yakınması ile başvuran dört yaşında kız olgu sunduk. Taşikardik olan hastanın elektrokardiyografisinde (EKG) V2, V3 ve aVF derivasyonlarında ters P dalgaları ile birlikte 0,16 sn P-R aralığı ve 0,28 sn R-P aralığı mevcuttu. Transtorasik ekokardiyografide sistolik fonksiyonları azalmış, geniş ve sferik sol ventrikül saptandı. Holter EKG'de kalp hızı 140-160/dk olan uzun R-P intervallli taşikardi konfirme edildi. Hastaya permanent resiprokan kavşak taşikardisine bağlı dilate kardiyomiyopati tanısı kondu ve hasta kateter ablasyonu ve flekainamid tedavisi ile başarılı bir şekilde tedavi edildi.

**Anahtar Sözcükler:** Elektrokardiyografi, çocuk, kateter ablasyonu

### Introduction

Permanent junctional reciprocating tachycardia (PJRT) is a rare form of orthodromic atrioventricular (AV) re-entry tachycardia with antegrade conduction through the AV node and retrograde conduction through a concealed accessory pathway which has very slow retrograde conduction. It is characterized by deeply inverted P waves in lead 2, 3, aVF and left precordial leads (1). It may be diagnosed incidentally, but mostly presents with dilated cardiomyopathy resulting from incessant tachycardia at a rate ranging from 120 to 250 beats/minute. PJRT is usually refractory to medical treatment, thus, early recognition and treatment with non-pharmacological methods, such as surgery or catheter ablation of the disease is essential (2). Herein, we present a four-year-old girl with PJRT-related

dilated cardiomyopathy who was successfully treated with catheter ablation and flecainide.

### Case

A 4-year-old girl was admitted to our hospital with the complaints of dyspnea, tachypnea, cough, excess sweating, and fatigue. She had been evaluated at another hospital where the chest X-ray revealed cardiomegaly. On admission to our hospital, physical examination revealed normal blood pressure (100/60 mmHg), tachycardia (180 beats/minute), tachypnea (50 breaths/minute), crackles at the basal areas of the lungs, hepatomegaly and a mild 1-2/6 systolic murmur. The electrocardiogram (ECG) showed inverted P waves in leads 2, 3 and aVF along with a P-R interval of 0.16 sec and an R-P interval of 0.28 sec (Figure 1). Transthoracic echocardiography revealed

**Address for Correspondence/Yazışma Adresi:** Şule Yıldırım

Çanakkale Onsekiz Mart University Faculty of Medicine, Department of Pediatrics, Çanakkale, Turkey

Phone: +90 505 828 07 07 E-mail: sule.yildirim@comu.edu.tr

**Received/Geliş Tarihi:** 14 April 2016 **Accepted/Kabul Tarihi:** 06 June 2016

The Turkish abstract was presented as a poster in "14<sup>th</sup> National Congress of Pediatric Cardiology and Cardiac Surgery" at 15-18 April 2015 in Denizli, Turkey.


© Copyright 2017 by The Medical Bulletin of Haseki Training and Research Hospital  
The Medical Bulletin of Haseki published by Galenos Yayınevi.

©Telif Hakkı 2017 Haseki Eğitim ve Araştırma Hastanesi  
Haseki Tıp Bülteni, Galenos Yayınevi tarafından basılmıştır.


an enlarged and spherical left ventricle with diminished systolic functions (ejection fraction: 27%, fractioned shortening: 13%) (Figures 2A and 2B). Biochemical and hematological parameters were within the normal limits. Viral markers were negative. Laboratory tests for inborn errors of metabolism were all negative. Cardiac magnetic resonance imaging was performed to detect possible myocarditis and it was normal. Digoxin, angiotensin-converting enzyme inhibitor and furosemide were started. Holter ECG confirmed long R-P tachycardia with a rate of 140-160 beats/minute. We used amiodarone and propafenone to achieve the sinus rhythm but we failed. Finally, the patient underwent radiofrequency ablation. The accessory pathway was interrupted successfully. No complication developed during the procedure. After two weeks of the procedure, Holter ECG was again positive for PJRT with a rate of 150 beats/minute. Before the second attempt for ablation, oral flecainide was started. After one week of flecainide, the ECG showed a sinus rhythm with a rate of 90-100 beats/minute and Holter was also normal. Additionally, one month after the flecainide treatment, transthoracic echocardiography demonstrated progressive improvement of the ventricular functions (ejection fraction was 57% and fractional shortening was 29%).

## Discussion

Tachycardia-induced dilated cardiomyopathy is a relatively rare but treatable type of cardiomyopathy. There



**Figure 1.** The electrocardiogram of the patient



**Figure 2.** A and B; transthoracic echocardiography images of the patient

are various tachyarrhythmias associated with tachycardia-induced cardiomyopathy, such as atrial fibrillation, atrial flutter, ectopic atrial tachycardia, incessant atrioventricular reciprocating tachycardia/PJRT, and ventricular tachycardia (3). Restoring the sinus rhythm or to slow the ventricular rate may result in an improvement in left ventricular functions.

Although the animal models have showed that rapid pacing produces marked depression of left ventricular ejection fraction, depressed cardiac output and increased systemic vascular resistance, the exact mechanism of tachycardia-induced cardiomyopathy is unclear (4). Depletion of myocardial energy stores and myocardial ischemia, abnormal calcium handling and beta adrenergic responsiveness and increased oxidative stress, and injury are the most studied mechanisms (5-8).

The manifestations of PJRT usually occur in childhood and may not be recognized until adult ages. The most important complication of PJRT is the development of tachycardia-induced cardiomyopathy (9). During PJRT-induced tachycardia the cardiac stimuli conduct antegrade through the AV node and return retrograde through a slowly conducting accessory pathway that is usually located near the ostium of the coronary sinus. It is characterized by deeply inverted P waves in lead 2, 3 and aVF. Our patient had the clinical signs and symptoms of congestive heart failure due to tachycardia-induced cardiomyopathy with specific ECG characteristics.

PJRT is usually resistant to medical treatment. Most of the patients with PJRT undergo various antiarrhythmic regimens to prevent tachycardia-induced cardiomyopathy. Most of the patients finally require catheter ablation procedures. Although there is a disagreement regarding the most useful therapeutic regimen, most of the studies support catheter ablation (9). Because the heart rate of infants and children with PJRT decreases with age, some reports suggest to postpone catheter ablation in older ages and to try medical regimens before ablation (10). Due to the resistance to medical regimen, we preferred ablation in our patient but after unsuccessful ablation, sinus rhythm was achieved with flecainide.

In conclusion, children with dilated cardiomyopathy should undergo a comprehensive evaluation of tachyarrhythmia such as PJRT. Medical regimens should be tried before ablation in small children and infants. If PJRT is considered refractory to medical regimens in the presence of ventricular dysfunction, catheter ablation should be preferred. Recognition and appropriate treatment of PJRT is important to prevent or reduce the complications such as tachycardia-induced cardiomyopathy.

## Ethics

Peer-review: Externally peer-reviewed.

### Authorship Contributions

Concept: Fatih Battal, Şule Yıldırım, Hakan Aylanç, Fatih Köksal Binnetoğlu, Nazan Kaymaz, Celal Akdeniz. Design: Fatih Battal, Şule Yıldırım, Hakan Aylanç, Fatih Köksal Binnetoğlu, Nazan Kaymaz, Celal Akdeniz. Data Collection or Processing: Fatih Battal, Şule Yıldırım, Hakan Aylanç, Fatih Köksal Binnetoğlu, Nazan Kaymaz, Celal Akdeniz. Analysis or Interpretation: Fatih Battal, Şule Yıldırım, Hakan Aylanç, Fatih Köksal Binnetoğlu, Nazan Kaymaz, Celal Akdeniz. Literature Search: Fatih Battal, Şule Yıldırım, Hakan Aylanç, Fatih Köksal Binnetoğlu, Nazan Kaymaz, Celal Akdeniz. Writing: Fatih Battal, Şule Yıldırım, Hakan Aylanç, Fatih Köksal Binnetoğlu, Nazan Kaymaz, Celal Akdeniz.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

### References

1. Aguinaga L, Primo J, Anguera I, et al. Long-term follow-up in patients with the permanent form of junctional reciprocating tachycardia treated with radiofrequency ablation. *Pacing Clin Electrophysiol* 1998;21:2073-8.
2. Cruz FE, Cheriex EC, Smeets JL, et al. Reversibility of tachycardia-induced cardiomyopathy after cure of incessant supraventricular tachycardia. *J Am Coll Cardiol* 1990;16:739-44.
3. Baman TS, Lange DC, Ilg KJ, et al. Relationship between burden of premature ventricular complexes and left ventricular function. *Heart Rhythm* 2010;7:865-9.
4. Spinale FG, Tomita M, Zellner JL, Cook JC, Crawford FA, Zile MR, et al. Collagen remodeling and changes in LV function during development and recovery from supraventricular tachycardia. *Am J Physiol* 1991;261:308-18.
5. Spinale FG, Clayton C, Tanaka R, et al. Myocardial Na<sup>+</sup>,K<sup>(+)</sup>-ATPase in tachycardia induced cardiomyopathy. *J Mol Cell Cardiol* 1992;24:277-94.
6. Perreault CL, Shannon RP, Komamura K, Vatner SF, Morgan JP, et al. Abnormalities in intracellular calcium regulation and contractile function in myocardium from dogs with pacing-induced heart failure. *J Clin Invest* 1992;89:932-8.
7. Marzo KP, Frey MJ, Wilson JR, et al. Beta-adrenergic receptor-G protein-adenylate cyclase complex in experimental canine congestive heart failure produced by rapid ventricular pacing. *Circ Res* 1991;69:1546-56.
8. Mihm MJ, Yu F, Carnes CA, et al. Impaired myofibrillar energetics and oxidative injury during human atrial fibrillation. *Circulation* 2001;104:174-80.
9. Kang KT, Potts JE, Radbill AE, et al. Permanent junctional reciprocating tachycardia in children: a multicenter experience. *Heart Rhythm* 2014;11:1426-32.
10. Semizel E, Ayabakan C, Ceviz N, Celiker A. Permanent form of junctional reciprocating tachycardia and tachycardia-induced cardiomyopathy treated by catheter ablation: a case report. *Turk J Pediatr* 2003;45:338-41.